

Patient Instructions following ULNAR NERVE DECOMPRESSION SURGERY

Dr. Noor Alolabi, Plastic and Reconstructive Surgeon



FOLLOW UP APPOINTMENT

Date: _____ **Location:** Fracture Clinic, Section D (Oakville Hospital)

SURGICAL SITE CARE AFTER SURGERY

- Keep the tensor bandage dry and clean for 4-7 days. You may then remove the outer dressing but keep the clear sticky bandage in place until your follow-up appointment.
- Move your elbow as much as the tensor bandage allows at least 5 times daily after surgery.
- Do not perform any heavy lifting with the hand for 6 weeks following surgery. You may use it for light activities such as getting dressed, eating, typing, etc.
- The cleaning solution used in surgery is pink. Do not be alarmed if your arm or fingertips are pink, it is not an infection.
- Your sutures are dissolvable and won't need to be removed

PAIN MANAGEMENT AFTER SURGERY

- The local anaesthetic (freezing) used will last for 4-8 hours and sometimes even longer. It is advised to take the prescribed or an over-the-counter pain medication once the tingling in the fingers starts to wear off. If you had a regional block and your whole arm is frozen, the anaesthetic can last 12-24 hours – do not be alarmed.
- Take the prescribed pain medication as needed for the first few days after surgery. You can supplement this with Advil if needed.
- Keep your hand elevated when sitting or lying down to minimize throbbing and swelling.

SCAR CONTROL AFTER SURGERY

- After 2 weeks from surgery, massage the scar multiple times a day for 2-3 months.
- Purchase "Silicone Gel" and apply over the scar for 2-3 months. This can be purchased from Dr. Alolabi's office.

DR. NOOR ALOLABI, MD, FRCSC
Plastic and Reconstructive Surgeon

560-2525 Old Bronte Road, Oakville, ON, L6M 4J2
Tel: (289) 386-3636 Fax: (289) 386-3536
www.drnalolabi.com